

SUPPLEMENTARY TABLE 1. Multivariable logistic regression analysis of possible predictors for silent brain infarct and cerebral microbleeds

	Silent brain infarct		Cerebral microbleeds	
	Adjusted OR (95% CI)	<i>p</i> -value	Adjusted OR (95% CI)	<i>p</i> -value
Age (years)	1.00 (0.99-1.02)	0.720	1.01 (0.98-1.03)	0.690
Sex (male)	0.99 (0.71-1.39)	0.961	0.52 (0.33-0.83)	0.006
Body mass index (kg/m ²)	0.98 (0.93-1.03)	0.489	0.95 (0.89-1.02)	0.156
Hypertension	1.06 (0.74-1.51)	0.772	1.14 (0.70-1.84)	0.606
Diabetes	1.22 (0.82-1.82)	0.337	1.30 (0.77-2.22)	0.326
Hyperlipidemia	0.98 (0.70-1.37)	0.902	1.01 (0.65-1.57)	0.964
Current smoking	0.82 (0.52-1.30)	0.398	1.65 (0.95-2.87)	0.078
White blood cell (x10 ³ /μL)	0.98 (0.89-1.08)	0.636	0.97 (0.86-1.11)	0.677
ESR (mm/hours)*	0.99 (0.89-1.09)	0.813	0.91 (0.79-1.04)	0.175
Anti-platelet agent	1.36 (0.89-2.08)	0.162	1.16 (0.64-2.10)	0.634

ESR; erythrocyte sedimentation rate.

*This variable was introduced as squared root scale.